



# CHILD GI PROGRAM\*

Support for Detoxification & Immune Function\*

- Start with ONE (1) spray of Biocidin® Throat Spray a day for the first three days. Then, add in ONE (1) drop of Biocidin® Liquid. Continue to increase, following the guidelines below.
- If the child experiences a Herxheimer/ cleansing reaction,<sup>†</sup> pause until symptoms disappear. Then, reintroduce the Biocidin® at the previous dose that did not produce symptoms.
- Only increase at a rate where the child is reaction free. It is not a race, but a journey to wellness.



## G.I. Detox™ Capsules

**CHILDREN:** Take ONE (1) capsule per 50 lbs of body weight.

**Take on an empty stomach 1-2 hours away from food, Biocidin®, other supplements, and/or medications.**

**BEGINNING DOSAGE:** ONE (1) capsule once a day per 50 lbs of body weight. Capsule can be opened and halved for children weighing less than 50 lbs.

**MAXIMUM DOSAGE:** Can increase to ONE (1) capsule twice a day per 50 lbs of body weight if child is having a Herxheimer/cleansing reaction<sup>†</sup> from the Biocidin®.

If the child is unable to swallow capsules, open the capsule and mix its content in water or applesauce.

This supplement is an essential part of the protocol to “mop up” and speed the removal of toxins.\* It should be used daily while on the Biocidin® protocol to avoid discomfort.\*



\*This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

## Biocidin® Liquid

**CHILDREN:** Start with the beginning dosage. If needed and well tolerated, gradually work up to the maximum dosage and maintain until symptoms improve.

**Take on an empty stomach ½ hour before a meal.**

**BEGINNING DOSAGE:** ONE (1) drop in a small amount of water once a day.  
Gargle and swallow.

If no Herxheimer/cleansing reactions<sup>†</sup> occur during the first three days, increase by one drop at a time until you reach the maximum dosage.

**MAXIMUM DOSAGE:** ONE (1) drop per 10 lbs of body weight in divided doses (morning and evening).



## Biocidin® Throat Spray

**CHILDREN:** Start with the beginning dosage. If needed and well tolerated, gradually work up to the maximum dosage and maintain until symptoms improve.

**Take on an empty stomach ½ hour before a meal.**

**BEGINNING DOSAGE:** ONE (1) spray once a day in the back of the throat.

If no Herxheimer/cleansing reactions<sup>†</sup> occur during the first three days, increase by one spray each day until you reach the maximum dosage.

**MAXIMUM DOSAGE:** FOUR (4) sprays twice a day.



<sup>†</sup>When toxins are released during cleansing and symptoms occur, it is called a “Herxheimer reaction” (herx). Symptoms can be different for each person and may include sore muscles and joints, fatigue, headaches, flu-like symptoms, changes in bowel habits, rashes, or other changes.

A herx is a result of detoxifying faster than your body’s detoxification pathways can handle. It’s a good sign that your cleansing is working. Just reduce the dosage and increase more slowly. Clinical evidence suggests that IF you are going to have a Herxheimer reaction (and it is rare), it will occur within three days. So, just increase the dosage every four days by ONE (1) until you reach the recommended dosage. Drink plenty of water and allow for extra rest when any signs of a herx occur.

**Remember, a sudden or severe reaction, such as an allergic reaction, is not a herx. Should a strong reaction occur, discontinue the use of any medication or supplement and contact your healthcare professional.**

Biocidin® and G.I. Detox™ are trademarks of Bio-Botanical Research Inc.